

# Comparing Agricultural Land-use Statistics from Different Sources: a Case Study in Greece\*

Vasiliki Benaki<sup>1</sup>, Luca Salvati<sup>2</sup>, Pier Matteo Barone<sup>3</sup>, Carlotta Ferrara<sup>3</sup>

<sup>1</sup> National Statistical Service of Greece, Pireos 46 & Eponiton, GR-18510 Pireas

<sup>2</sup> Italian Agricultural Research Council, Centre for the Study of Soil-Plant Interactions, Via della Navicella 2-4, I-00186 Rome

<sup>3</sup> Department of Physics "E. Amaldi", University of Roma Tre, Via della Vasca Navale, 84, 00146 Rome, Italy

<sup>1</sup> vasbenaki@statistics.gr

<sup>2</sup> luca.salvati@entecra.it (corresponding author)

<sup>3</sup> ferrara.fis@uniroma3.it

\*A preliminary draft of this contribution was presented in the 'First Conference of the Hellenic Association of Information and Communication Technology in Agriculture, Food and Environment' (Athens, 2002).

**Abstract** - This paper illustrates a tool integrating the Farm Structure Survey (FSS) and an improved version of CORINE Land Cover (CLC) map as a contribution to produce reliable land-use statistics at national and regional scale. To achieve compatibility between census and CLC the present tool takes into account the FSS nomenclature and definitions, and reorganizes the 44 classes of the original CLC into 16 general classes that meet the needs of the Land Use/Cover statistics in Greece. To compare the respective surface areas of the related classes and to provide the appropriate links between certain classes of the two nomenclatures four Greek regions are used in the pilot study: Kriti, and the three prefectures of Makedonia. The linkage between the two databases shows the existing differences between the administrative areas. The developed tool is able to relate data from different sources, and to display on a map, the combined spatial statistical data along with the geographical information of the area. Thus, although the new CLC seems to provide a good mapping base, the imposed minimum mapping unit of 25 ha results in an overall underestimation of the diversity of agricultural land-uses, something which is particularly important in the case of Greece for which the average size of the farm holdings is less than 5 hectares.

**Keywords** - Farm Survey Structure, Corine Land Cover, Landscape diversity, Greece

## 1. Introduction

Agricultural activities are more and more combined with other activities such as environmental protection, maintaining the landscape, forestry, preserving recreational and tourist areas as well as

small scale of agricultural products. Thus, there is a strong need for statistical data on rural population, and particularly, on landscape and land-use. Most of the statistical data used for policy purposes are related to populations, activities, features and other events, which are by their nature, spatial in form. The management, the process and the display of statistical data associated with spatial locations that vary geographically is mainly a spatial process. In agricultural terms, the management of agricultural resources is increasingly complex as conservation and environmental concerns play an expanding role for making conclusions. In this respect, Geographic Information Systems are needed in the production of census maps, for dealing with census logistics, for monitoring census activities, and for data dissemination (Deichmann, 1997). A wide range of spatial analysis methods has been developed for carrying out data transformations between different spatial structures. These methods help to present the data in a more meaningful and consistent manner and enable different data sets, based on different geographical units, to be brought together and overlaid. They also facilitate the spatial analysis of the statistical data required in the development of some more reliable indicators for the determination of the state and quality of the environment, able to measure the effect of the agricultural economy, across regions and countries. The use of indicators as an aid to policy decision-making in the agri-environmental context is a relatively recent phenomenon and still a developing field; however, indicators are perceived to have considerable potential as policy tools. Most policy makers

concerned with agri-environmental issues at the national level are confronted with fragmented information and it is accordingly difficult to use the information in a way that effectively contributes to policy decision-making. An unavoidable step in the assessment of agricultural policies and of their impact on the countryside and landscapes is the study of spatial units that constitute the underlying structure of these territories. Most statistical data in EU, by means of the Farm Structure Survey (FSS) data, is organized and presented on the basis of NUTS (Nomenclature des Unites Territoriales Statistiques) system, to provide a single, uniform breakdown of a country. Nevertheless, these units are geographical areas that may vary substantially not only in the sizes and shapes, but also over the time. In addition, this geographical level is not appropriate to carry out certain environmental studies. The need of spatial analysis and of the production of environmental indicators requires delineation of the land use data according to their natural depiction on a geographical map, beyond the administrative distribution. As a result, NUTS system cannot be applied in its present form to units that are more relevant from a geographical point of view, such as drainage areas, landscape units, biotopes, etc. This study illustrates the development of a tool interface between statistical and geographical databases by comparing Farm Structure Survey (FSS) and CORINE Land Cover (CLC) land-use figures. As a first step, the spatial disaggregation of the FSS data into an accurate geographical level requires an interface between the two nomenclatures. To reallocate the FSS data into sustainable areas a question arises of how the digital CLC map could be used to describe agro-environmental statistical structures. Note that CLC has so far been focused on land cover, rather than land use and it has been carried out once. As a result it cannot be applied to show trends. However, different countries carried it out in different years, over the period 1985- 1995. Plans already exist to upgrade CLC based on the IMAGE 2000 image data set provided by the JRC. The result is that some of the indicators based on CLC show only a snapshot rather than a trend in land-use. The developed interface is able to display on a map, accurately, the combined spatial descriptive statistical data along with the geographic information of an area of interest. Thus, the user is able to relate the FSS and the CLC data in order to find the best matching. The developed interface is able to query a database, aggregate / disaggregate the data and plot the results

on a map. The comparison requires to determine the aggregation level of the classes for which the correspondence has already been set and to validate the result by comparing the respective surface areas of the related classes. After the reclassification of the above data, common classes are created and presented on a map using an embedded GIS environment. To test the interface and provide the appropriate links between certain classes of the two databases the three regions of Makedonia and the region of the island of Kriti have been chosen. The statistical data used has been provided by the Basic FSS of 1999/2000 (Census of Agricultural for Livestock breeding or simply Agricultural Census). However, to achieve compatibility between census and photo-interpretation a recently developed, improved version of the CLC geographical database has been used. The new CLC takes into account the FSS nomenclature and definitions and has provides much better acquisition period (Landsat-TM 1998 to 1999) which is the same with the census reference period (1998 to 1999). The linkage between the two databases shows the existing differences between the administrative areas of the pilot regions. The structure of the paper is as follows: chapter 2 describes the main characteristics of the FSS nomenclature, particularly addressed in the case of Greece. Chapter 3 addresses the modified CLC geographical nomenclature providing the new classification scheme. Also, in this section, the original CLC nomenclature is discussed briefly. Chapter 4 discusses the linkage between the two nomenclatures and the way it has been achieved by means of application development. Chapter 5 illustrates the results starting from the comparison of the related nomenclatures and finally, in the last section the conclusions of this work are presented.

## **2. The FSS database**

### **2.1 Main issues**

The effective and balanced implementation of the reformed Common Agricultural Policy requires detailed objective, quantitative data of the structure and performance of the agricultural, rural and environmental sectors. In this context the development of the structure of the agricultural holdings allows analysis of the agricultural sector and its impact to other sectors as the rural sector and the environment. The FSS is the main source to provide data on various characteristics relating to agricultural holdings, on a regular basis. These data refer to the number and size distribution of the

agricultural holdings by type of enterprise, as well as to the land improvements, crop and livestock rotations and farm practices (machinery, equipment etc.). They also refer to other structural data such as the educational level of farmer and farm labor inputs, the legal status of holder including tenure arrangements and finally other social demographic characteristics of holders. The FSS data are collected on a regular basis by the Member States and are forwarded to Eurostat, which stores them in the Eurofarm database. In order to harmonize information at the Community level, legal frameworks (Regulation and Decisions) define the methodological framework and the contents of the FSS questionnaires. Table 1 shows the FSS nomenclature, which distinguishes the detailed agricultural land use classes.

## 2.2 Methodological issues of the FSS in Greece

The FSS is carried out in Greece within the framework of the Community Program for the 'Statistical Surveys in the Agricultural Sector'. All the specifications and terms are defined precisely by the Regulation 571/88 as amended by the Regulation 2467/96 and the related Decisions of the Council of the E.U. The FSS is intended to collect statistical data on the structure of agricultural and livestock holdings and the employment of the population on them. The data make it possible, besides the classical tabulation of the results, to generate tables, which show the economic size and orientation of the farms (typology). In particular, the Greek FSS system aims to collect data relating to:

- The number of agricultural and livestock holdings in the country, at national, regional and local level.
- The geographical position of the holdings.
- The legal status and management of the holding.
- The agricultural training of its owner.
- The keeping of account books.
- The land uses (arable crops, permanent crops, kitchen gardens, permanent pasture-meadows and rough grazing and other areas).
- The type of ownership of the utilized agricultural area.
- The number of fields constituting the total

utilized agricultural area.

- Successive crops, combined crops, irrigated crops, etc..
- Livestock raised on the farm..
- Agricultural machinery and milking equipment used.
- Employment of members of the farm owner's family.
- Employment of family members in other gainful activities besides agriculture.
- Employment of permanent, seasonal and other workers.

Sample FSS is carried out every two years, in the years ending with an odd number. The National Statistical Service of Greece (NSSG) carried out the first sample survey of the Structure of Agricultural and Livestock breeding in 1966/67, when Greece was still an associated member of the EU. The next sample survey took place in 1977/78. After the accession of the country into the EU further surveys were carried out in 1983, 1985, 1987, 1989, 1993, 1995 and 1997 i.e. every two years. Every ten years an exhaustive survey (Basic FSS or Agricultural Census) is carried out. The first Agricultural Census conducted in 1950, after the Second World War. Agricultural Census of 1991 was the last census carried out at the same time with the General Censuses for population, households, agriculture etc. However, Agricultural Census in 2000 was the first census carried out before the General Population Census dated 2001 and it was based on the Farm Register. The reference period for the data collected on crops and employment is from October 1st of year t-1, to 30 September of year t, i.e. the survey year. Exceptions to this are a farm's livestock and machinery, questions relating to which have a reference date of 30 September in the year t, for the machinery and 1st November for the livestock. The statistical unit for the FSS is defined as an agricultural or livestock holding<sup>4</sup> which during the reference period comprises at least one of the following:

- at least 0.1 ha of utilized agricultural area or at least 0.05 ha of greenhouse area, regardless of its own ship and location, or
- at least one cow, or

- at least two other large animals (oxen, buffaloes, horses, etc.), or
- at least five small animals (sheep, goats, pigs), or
- at least 50 poultry birds, or
- at least 20 beehives.

The FSS was carried out by filling in a special questionnaire after interviewing the owner of the agricultural or livestock breeding farm. The sample survey is carried out by applying the method of multi-stage stratified area sampling. In the most recent Agricultural Census in Greece the Basic FSS covered all agricultural and livestock holdings in the country (nearly 814.000 holdings).

### 3. Materials and methods

#### 3.1 The CLC geographical database

CORINE (Co-ORDination on INformation of the Environment) Land Cover (CLC) is a geographic land cover/land use database encompassing most of the countries of the European Community, with aim to gather information associated with the environment on certain priority topics. It describes land cover (and partly land use) according to a nomenclature of 44 classes organized hierarchically in three levels (Dueker, 1979). CLC was elaborated based on the visual interpretation of satellite images (*Spot*, *Landsat TM* and *MSS*). The smallest surfaces mapped (mapping units) correspond to 25 hectares. Linear features less than 100m in width are not considered. The scale of the output product was fixed at 1:100.000. Thus, the location precision of the CLC database is 100m. Although its exploitation is just starting, it offers the potential for a wide array of uses. It can be used on its own for simple cartographic or statistical presentations and as a base for European-wide landscape analyses or more generally in combination with other data sets.

#### 3.2 The CLC database of Greece

The CLC database has been developed in Greece in order to cover the needs of land use/cover statistics as far as the distribution of the total area of Greece in the basic categories of land use is concerned. These statistics are included in the preparatory work carried out in the context of every Agricultural Censuses. The aim is to prepare the census and to obtain data covering all the territory of Greece. Until Agricultural Census of 1991, this work was done by completing seven (7) months before the Census a 'pre-census

questionnaire of total land area in the municipality or commune', using estimates by the municipal or communal working parties set up for the census and with the help of local agronomists. To facilitate completion of the pre-census questionnaire, these groups had at their disposal the land distribution data from the previous census, as well as other auxiliary data held by the municipality or commune, such as land registers, land distribution tables, etc. Land was divided up into seven basic categories of use:

- Cultivated areas and fallow land resting fallow for 1 to 5 years.
- Communal or municipal pasture land.
- Other pasture land (owned by privates, State, monasteries, etc.)
- Forests
- Areas under water (lakes, marshes, seashores, river beds)
- Build-up areas (buildings, courtyards and roads, squares etc.)
- Other areas (e.g. rocky areas, mines, etc.).

Note that the pre-census questionnaire was the only data source covering also the state-owned land, which is mostly, forest and pastures. Nevertheless, since the agricultural census is carried out by interviews of farmers it concerns only private lands that are somehow agriculturally used.

In the light of the recent developments concerning land use statistics, NSSG decided to use an up-to-date methodology using GIS techniques in order to produce more objective information on this sector. Therefore, the use of spatial analysis is required. Spatial analysis of the information to be recorded is realized by determining the area of the minimum recorded surface, which is taken according to the proposed nomenclature, the methodology of use/cover definition, the requirements of 1:100.000 scale and the user needs. The method with which the theme information drawn up, is the comparative photo-interpretation of the satellite data collected in 1998-99 in relation to those of the time period 1997-98 used for the creation of the CLC database in Greece. The digital photo-interpretation of the new satellite data is made using image processing software and other data such as those from land recordings. The recording planning and the use of the

data from the field works are also defining the reliability of the specific photo-interpretation.

The new CLC database is properly generalized as reference data and harmonized with the FSS nomenclature, by means of characteristics and definitions, linkage of the two databases to meet the needs of the NSSG. Thus, the distribution of the main land uses in Greece has been reorganized into the following sixteen (16) classes:

- Artificial surfaces
  1. Urban fabric
  2. Industrial and commercial units
  3. Transport units
  4. Mine, dump and construction sites
  5. Artificial, non-agricultural vegetated areas sport and cultural activity sites
- Agricultural areas
  6. Arable land
  7. Permanent crops
  8. Pastures
  9. Heterogeneous agricultural areas
- Forest and semi-natural areas
  10. Forests
    11. Transitional woodland /shrub
    12. Shrub and/or herbaceous vegetation associations
    13. Open spaces with little or no vegetation
- Surfaces under water
  14. Inland water
  15. Inland wetlands
  16. Coastal wetlands

The new CLC geographical database for the country's area has numerous advantages, the most important of which are the following:

- It provides a land use/cover map covering all Greece for 16 categories, compiled with the seven land use classes in the above mentioned pre-census questionnaire of the NSSG.

- The new geographical database takes into account the FSS nomenclature and definitions.
- It enables comparability between the two sources of information, namely census versus photo-interpretation. In the case of Greece the acquisition period of the data is spread over 2 years for both, the CLC (Landsat-TM 1998 to 1999) and the FSS 1999/2000, (reference year the 1998-1999 crop year).
- It enables the integration of the chrono-geographical co-ordinates of the satellite images sources of CLC. This will help in the identification of districts for which CLC's image interpretation is one year apart (minus or plus) from the census year (1990 or 2000, respectively). In addition, using the intermediate FSS data that correspond closely to the date of the satellite image it will be possible to mitigate the effect of time.

#### 4. Linking the two databases

As it is well known, data collection methods are optimized for a particular need and therefore the resultant data structures are not usually readily comparable in a cross-sectional study. Thus, although a particular census may be analyzed in detail comparing censuses with each other have been proved problematic since they may use different administrative units, or they may use the same unit system, which includes many boundary changes that make the comparison difficult. To the best of our knowledge, three types of data incompatibilities have been distinguished so far (Frank, 1999; Gregory, 2000) and will be described, briefly, below.

##### 4.1 Differences in Data Models

Raster and vector data models are the GIS approaches for the spatial presentation of natural vegetation, the forest area and generally the development of land use. In a raster data model, a uniform grid, each cell of which is assigned a unique descriptor depending on the coordinate system used, represents space. Raster models can be directly imported into the software and immediately become available for use (Burrough, 1986). They are well suited for the representation of remotely sensed digital data and are commonly employed in the environmental sciences. In contrast, in a vector data model, the spatial data is based on geometric shapes of points, lines, and polygons. This model is object-oriented and is based on the coordinate system used.

Vector GIS knows where the spatial feature (line, point, polygon) exists, as well as the relationship with the other features. Vector data models are particularly suited for the representation of linear data features like roads, or clearly delineated areas, such as, property lines and city limits. After the representation of the spatial features, their associated properties must be specified in a separate database. For simultaneous use of data from both, raster and vector models a conversion of one data set to the respective model of the other data set needs to be performed. Data conversions, however, are often ambiguous and typically result in a loss of information (Maffini et al., 1987). It is difficult, for example, to derive the best fitting vector representation from a given raster grid. Furthermore, it is likely that a set of transformations from vector to raster and back to vector will result in a target feature whose shape differs from the original source feature. These transformation functions may not be accessible to the end user (Ehlers et al., 1991, Maffini et al., 1987). The data transformation from analog paper maps and tabulations to digital data falls into the same category. Loss of data, spatial inaccuracy and error is introduced by conversion techniques like scanning, digitizing, rasterization, vectorization and manual data input (Goodchild, et al., 1989). Digital data creation is also extremely time consuming.

#### 4.1.1 *Inconsistencies of areal units*

Comparing census data with other data sources of some specific area of interest may not be the same, either because of boundaries changes over the time or because of the different definition of administrative/areal units used for the data collection (Xie et al., 1995). Two key issues need to be addressed in terms of areal unit comparability. One is related to data integration and map overlay (non-matching areal units). The other is related to data analysis and statistical comparability of areal units of different sizes and shapes (modifiable areal unit problem).

#### 4.1.2 *Non-matching Areal Units*

Integrated analysis of spatial and attribute data is based on map overlay operations. Non-matching areal units require a transformation of data from one system of areal units to another in which data values are apportioned to the newly created spatial units. Then, the newly created zones allow data overlay and analysis. These transformations are known as "areal interpolation".

#### 4.1.3 *Modifiable Areal Unit Problem (MAUP)*

Generally, the statistical data, whose distribution and characteristics are not well known, are presented by an appropriate aggregated variable of some higher class. In addition, censuses base their statistics on well-defined areal units that tend to vary in size and shape leading to inconsistent and misleading statistical results. This is known as a Modifiable Areal Unit Problem (MAUP) (Openshaw, 1984). A possible approach to face this problem would be the reaggregation of the available data into homogeneous subunits and the increase the spatial detail using ancillary land cover data in order to display the census data on a map (Yuan et al., 1997).

#### 4.1.4 *Temporal incompatibilities*

Data collection of land cover data and monitoring of physical changes relies on remote sensing via aircraft or satellites. Coverage cycles, for example, for the different LandSat orbiters (Lillesand, et al., 1994) range from 8-14 days. This data density, however, may be deceptive since data for certain regions is usually available for much fewer dates due to the fact that frequent cloud cover prohibits data collection. In the case of agricultural cultivations, the above problem has to be considered in more detail since the cultivations are usually visible on specific periods over the year.

## 4.2 **Technical procedure**

To describe the methodology adopted in the investigated issue, one has to take into account the non-matching areal units and the MAUP problems mentioned in section 4.1. The temporal incompatibilities problem and the procedure of matching the data points by non-matching due to collection cycles will not be considered because in our case, the data used has been interpreted by an independent intuitional organization. The Non-matching Areal Units problem arises due to the following reasons.

The different boundaries definition of the administrative units that have been used in the Hellenic censuses of the year 1991 and 2000. To solve the problem there is the need to transform the data between different spatial structures. As transformation may be described the process of aggregation and disaggregation within nested and non-nested neighbor polygons. To overlay the data together the conceptual model has been designed containing and maintaining all polygons and the

related geometric data (lines, nodes etc), representing the areal units. To link the descriptive and the spatial information, the data of the geographic area of interest is broken down into smaller parts in order to determine the field that identifies the specific entity ('AreaCode') in order to be used as a reference key to the GIS. A set of spatial queries also has been developed for carrying out the transformations.

Moreover, the different geodetic datums used for the presentation of statistical and for the ancillary geographical data. To use the ancillary data along with the other geographical data a target datum has to be selected as the reference datum for all data and an automated procedure has to be developed to convert the data between the source and the selected target geodetic datum.

To automate both the transformation between different definitions of administrative units and the conversion between different geodetic datums, an object has been designed, called "Geo-Object". This Object can be used as the basic map layer on any similar application. The MAUP problem is faced using ancillary geographic data (Flowerdew et al., 1991) such as contour lines, lines representing rivers or polygons representing lakes. This methodology permits the synthesis of geographical data along with the studied statistical data and allows the combination of different scenarios in order to simulate the plotting of the descriptive data containing quantitative and areal information on a map. For validation and / or prediction purposes, the results are compared visually with other spatial quantitative information or sampling data presented on thematic maps.

A database entity object provides the connection between the conceptual model and the input quantitative data. It has only one method that is used to insert a new 'AreaCode' into the conceptual model and finally it connects the new inserted 'AreaCode' with the appropriate areal information. A tab-delimited text file contains some quantitative information in a country level. To link this text with the appropriate geographic feature in the GIS environment, the entity "Details" of the conceptual model must be updated with the 'AreaCode' of countries that this text file contains. This can be done easily by using the properties and methods of the above described class of objects.

The advantage of the described methodology is the capability to combine quantitative data from different sources, and to compare them with the available spatial features concerning the distribution

of similar quantitative data (thematic map) of the same area of interest, into an integrated geographical environment. This environment can contain more geographical features than the ordinary thematic maps such as contour lines, roads, rivers, airport, etc.

This can also be helpful for the methods that have been developed to solve the problem of geographic missing values. The precision of those methods is depended on the availability and quantity of historical data. Using this methodology, it is obvious that the integration of all types of GIS data with quantitative data available from other sources is crucial for someone to decide about the data correctness.

### 4.3 Study site application

As it has been pointed out, the linkage of the two nomenclatures, by means of the FSS the CLC databases require computer-based application software able to display maps and descriptive data in a tabular form. This has been achieved using geographical information from CLC database linked with tabular information of the multi-dimensional tables of FSS (Table 2). The user becomes part of the GIS without the necessity of specific skills and intimate knowledge of the data used. The application consists of the following parts:

- A relational database
- The class of objects for data manipulation
- The class of objects for GIS manipulation
- The main body of the application software containing the above items along with the functions required by the end user.

To begin with, a step-by-step analysis of the software design is required. However, for the purpose of this research it is assumed that the pilot area is already known. Then, the appropriate design steps are as follows:

1. On the CLC's geographic layer of the area of interest we add the remaining geographic characteristics (contour lines, roads, cities, lakes, rivers etc.). This will help in the understanding the exact location of the CLC data.
2. From the FSS database we select only the themes, which associated with agriculture products. The data selected is at prefecture

- level, in thousands of hectares of agriculture products, reported in 2000 census.
3. The data provided by the FSS and the CLC databases is studied in order to develop the entity relationship model, and then the database system of the application.
  4. CLC data is stored in some database tables of the application, using especially developed programs, while NC's data were stored manually. NC provides also the appropriate DLLs in order to develop programs for automated data transfer.
  5. We pointed out the appropriate functions and queries, and we developed object classes to satisfy the requirements for uniformity at both, user and developer levels.
  6. We developed an application in which are used the RDBMS, the GIS and the pre-mentioned object classes. The basic capabilities offered by this application are the following:
    - Ability to compose (aggregate) a new FSS theme by selecting one or more CLC classes, and vice versa.
    - Ability to decompose (disaggregate) an existing FSS theme to one or more CLC classes, and vice versa.
    - Ability to correspond (relate) the new FSS themes to CLC classes.
    - Ability to classify (sort) the results either by date, or by county (region), or by CLC class.

Ability to observe the results plotted on the map and to classify these by geographic characteristics, such as allocation of the selected growth by elevation.

## 5. Results

Table 2 presents the linkage between the 2000 FSS and the new CLC nomenclatures. Although the new CLC nomenclature has been harmonized with the FSS nomenclature there are still some problems related to the two different methodologies. The analysis of the above problems has been carried out throughout of a comparison between the respective areas of the related classes, and has been allowed to make proposals for a future work. The available data from the 2000 FSS has been based at Municipality/Commune level (NUTS V or LAU-1

level), whereas the data has been drawn from the new CLC at the district level (NUTS III). The data of two databases has been compared in a pilot study of four regions of Greece at a district level (NUTS III). The comparison shows large deviations in the agricultural areas. Generally, the examined agricultural areas in new CLC are greater than the corresponding agricultural areas in the 2000 FSS. The problem of large deviations is caused mainly because of the difficulties in correlating the pastures areas between the two databases, whereas the differences of the arable areas and the areas under permanent crops are related to the different methodologies.

The results found so far are presented in Tables 3 to 5. Table 3 presents the differences (%) in arable areas, areas under permanent crops, and cultivated areas (aggregation of D+E), as they recorded in the districts (NUTS III) of the examined regions, between the two nomenclatures. Positive sign is in favor of the new CLC nomenclature, whereas negative sign is in favor of the FSS nomenclature. Note that the actual differences in the above classes are not as high as they are in the remaining classes, namely pastures and meadows (Table 4), heterogeneous areas and agricultural areas (Table 5). To facilitate the comparison for the last cases the actual values are presented.

As it may be observed (Table 3) the above differences (%) in the regions (NUTS II) are generally smaller from the corresponding inter-regional ones (district level; NUTS III). This is due to the fact that the mapping unit of 25 ha in the new CLC is not able to identify parcels of smaller size. This is the case of Greece, in which the average holding size is around 4.5 ha and the average parcel size is around 0.7 ha. An additional reason is that in FSS all the holdings are recorded at the place of residence of the holder (natural person) or headquarter (legal person) of the holding. In the following some preliminary comparison of these results are summarized:

- *Arable areas*

Region comparison shows that the difference for the region of Kriti is about 66% in favor of the FSS nomenclature. However, the differences in the regions of Makedonia are not as high (at most 33%) and are in favor of the new CLC nomenclature (Table 3). Generally, the differences in the arable crop areas are moderate and are in favor either of the FSS nomenclature or of the new CLC nomenclature (NUTS III level). Interesting to note that in some



districts of the regions of Kentriki and Dytiki Makedonia the results are almost the same.

- *Areas under permanent crops*

In general terms the situation is opposite of the one described in the arable crop areas. As it may be observed from Table 3 in the region of Kriti the differences (%) between the two nomenclatures are very small (about 6%). In the regions of Makedonia these differences (%) are moderate (at most 61%) and are in favor of the FSS nomenclature (NUTS II). Furthermore, in the districts of some regions these differences are substantial and/or in the opposite direction (e.g. Evros, Rodopi).

- *Pastures and meadows*

The total areas of pastures and meadows are generally larger in the new CLC than the corresponding areas recorded by the FSS. In all regions (NUTS II) the differences are very high. In the region of Kriti, the two districts of Rethimno and Chania the recorded areas in the new CLC are smaller than the corresponding areas of FSS. This is because only the private areas are recorded in the FSS, whereas all pastures (such as state-owned pastures, private pastures, etc.) are recorded in the new CLC.

- *Heterogeneous areas*

FSS and new CLC present very high actual differences in the class of the heterogeneous areas. Even the two nomenclatures are harmonized there is still a methodological problem of how to relate the two nomenclatures. In particular, in the FSS the survey unit is the agricultural holding, which comprises of at least 0,1 ha. Therefore the heterogeneity (combined crops) of these areas is referred to this small area. In new CLC the heterogeneity is examined within the mapping unit of 25 ha. Under these circumstances a polygon in the new CLC that includes different parcels of a single crop is recorded as heterogeneous area, whereas in the FSS the corresponding parcels are recorded as single crops.

- *Agricultural areas*

All the Agricultural Areas (AA=D+G+F) resulting from the new CLC nomenclature show larger values than the corresponding areas in the FSS nomenclature, particularly in the districts. The differences are generally high with exception of two districts of the region of Kriti. As it has been pointed out previously, the large deviations observed between

the agricultural areas as they recorded in the new CLC and the FSS are due to the large deviations in the pastures.

- *Cultivated areas*

Given the problems of the large deviations in the total agricultural areas that are caused mainly from the pastures, the aggregation of the arable areas and the areas under permanent crops into the new class of "Cultivated areas" shows that the differences presented in this class are not significant.

## 6. Discussion

This study has been based on the provisional data of the 2000 FSS and the new CLC databases and it may be considered as a first step in the direction of present georeference statistical data. The difficulties appeared in the linkage of the two databases can be generally explained from the following points.

The different methodology used as far as the data collection methods and the coverage are concerned. In particular, the FSS is a census using as a reference unit the farm, whereas the new CLC is based on photo—intepretation of the whole area of the country using as a reference unit the mapping unit of at least 25 ha. In addition, CLC has so far been focused on land cover, rather than land use. The minimum size of 25 ha of CLC mapping units presents the difficulty of identifying parcels of smaller size. Thus, a number of non-agricultural areas are classified as agricultural whereas they are only partially agriculture. This is a common problem in areas with forest and olive-trees. Besides, areas classified as non-agricultural areas in CLC may include part of an agricultural area. This explains a number of differences within the agricultural classes. For example, part of meadows or permanent crops can be included in areas with arable crops and conversely.

Despite the harmonization between the new CLC and FSS nomenclatures there are still problems as far as pastures and heterogeneous areas are concerned. In the new CLC, the non-agricultural classes defined by the codes 11, 12, and 13 ("Transitional woodland/shrub", "areas with mixed shrub/grass vegetation" and "areas with little or no vegetation" respectively) may include surfaces classified as "permanent meadows and pastures" in the FSS. Furthermore the FSS does not record the state-owned meadows, which in the new CLC are recorded under the code 8 ("areas under meadow or

pastures"). The special features of Greek agriculture that is marked by the diversity of the holdings in terms of area of production (mixed holdings), the small size of the holdings (average size 4,5 ha), the fragmentation of their area (6 parcels approximately per holding and average parcel size of 0,7 ha). In quite a number of cases the parcels of the same holding are normally located far away from the farmhouse or from the headquarter, but they are recorded at the place of the farmhouse or the headquarter (by definition).

## 7. Conclusion

The work presented so far is a pilot study merging, by means of a software tool, the statistical data, available at the administrative level, with the geo-referenced land cover in order to identify and explain the most significant differences encountered between the aggregates of agricultural land cover classes. This has been achieved with the use of the 2000 FSS and the new CLC databases already under development in Greece. The new CLC seems to provide a good mapping base for Greece, which could be improved further by using suitable satellite images that are able to produce scaled maps of at least 1:50000. Note that the imposed minimum mapping unit of 25 ha results in an overall underestimation of the diversity of landscapes something, which is particularly important in the case of Greece for which the average size of the holdings is 4,5 ha. Apart of CLC, additional sources may be used providing detailed complementary information, such as aerial ortho-photographs, the cadastral map of Greece, IACS (Integrated Administrative Control System), MARS (Monitor Agriculture with Remote Sensing), NATURA2000 database, or other ongoing analysis of the European landscape.

When the final data from the remaining regions of Greece will be available a quality analysis of the two databases will be carried out and a finer level of nomenclature will be examined. This will allow final conclusions to be drawn and further actions to be taken in the future. Future research is to continue improving the idea of interoperable geo-object by adding methods and properties for uncertainty manipulation and to investigate requirements of GIS in a fuzzy object data model. Our final objective is to embody in the Geo-Object, the ability to generate and visualize transitions from one state to another, using the rules of an expert spatiotemporal system.

## References

- [1] A. Frank (19-21 June 1999), Tracing socioeconomic pattern of urban development, Proceedings of Geoinformatics'99 conference, Ann Arbor, Michigan.
- [2] Burrough P.A (1986), Principles of Geographic Information Systems, 1st ed, pp.336, Oxford University Press, New York.
- [3] Deichmann U. (6-10 December 1997), Geographical information systems in the census process —Technology options, costs and benefits, Workshop on Strategies for the 2000 Round of Population and Housing Censuses in the ESCWA Region Cairo.
- [4] Dragicevic V. and Marceau D. (2000), An application of fuzzy logic reasoning for GIS temporal modeling of dynamic processes, Fuzzy sets and Systems 113, Elsevier, pp.69-80.
- [5] Dueker, K.J. (1979), Land Resource Information Systems: A Review of Fifteen Years Experience, Geo-Processing, 1, pp.105-128.
- [6] Ehlers, M., Greenle, D., Smith, T. and Star, J. (1991), Integration of Remote Sensing and GIS: Data and Data Access, Photogrammetric Engineering & Remote Sensing 57(6), pp.669-675.
- [7] Flowerdew, Robin and Mick Green (1994), Areal Interpolation and Types of Data. In Spatial Analysis and GIS edited by Stewart Fortheringham and Peter Rogerson. London, Bristol: Taylor & Francis Ltd.
- [8] Goodchild, M. and Gopal S. (eds), (1989), Accuracy of Spatial Databases, London: Taylor and Francis.
- [9] Ian N Gregory (23 - 25 August 2000), An evaluation of the accuracy of the areal interpolation of data for the analysis of long-term change in England and Wales, Proceedings of the 5th International Conference on GeoComputation University of Greenwich, United Kingdom.
- [10] Lillesand, T.M. and Kiefer, R.W. (1994), Remote Sensing and Image Interpretation (3rd ed.), New York: Wiley.

- [11] Maffini and Guilio (1987), Raster versus Vector Data Encoding and Handling: A Commentary. *Photogrammetric Engineering & Remote Sensing* 53 (10), pp.1397- 1398.
- [12] Openshaw, S. (1984), *The Modifiable Areal Unit Problem*, Norwich: Geo Books.
- [13] Ozemoy, V.M., Smith, D.R., and Sicherman, A. (1981), Evaluating Computerized Geographic Information Systems Using Decision Analysis, *Interfaces*, 11, pp.92-98.
- [14] Parent C., Spaccapietra S., Zimanyi E., 'Conceptual Modeling for Federated GIS over the Web'. *Ecole Polytechnique Federale de Lausanne — Laboratoire de Bases de Données*, 1.
- [15] Sambrakos M., Fillis I., Tsiligiridis T. (2001), Integration of Spatial Descriptive Statistical Data and Geographic Information, *Proceedings of 8th Panhellenic Conference on Informatics*, paper 91C, Nicosia, Cyprus.
- [16] Xie and Yichun (1995), The Overlaid Network Algorithms for Areal Interpolation Problem, *Computers, Environment and Urban Systems*, 19 (4), pp.287-306.
- [17] Yuan, Y., Smith, R.M. and Limp, W.F. (1997), Remodeling Census Population with Spatial Information from Landsat TM Imagery, *Computers, Environment and Urban Systems*, 21(3/4), pp.245-258.

**Table 1.** Classification of land use in the 2000 FSS nomenclature.

		D01	Common wheat and spelt
D:	D01-D08: CEREALS	D02	Durum wheat
ARABLE LAND		D03	Rye
		D04	Barley
		D05	Oats
		D06	Grain maize
		D07	Rice
		D08	Other cereals
		D09C	Pulses-fodder peas
	D09: DRIED PULSES	DO9D	Pulses-fodder field beans
		DO9E	Pulses-other than fodder peas and field beans
		D10	Potatoes
	D10-D12: ROOT CROPS	D11	Sugar beets
		D12	Fodder roots and brassicas
		D13A	Tobacco
	D13: INDUSTRIAL PLANTS	D13B	Hops
		D13C	Cotton
		D13D	Other industrial plants
		D13D1	Other oil seeds or fibre plants
		D13D1A	Rape and turniprape
		D13D1B	Sunflower
		D13D1C	Soya
		D13D1D	Other oil seeds or fibre plants-others
		D13D2	Aromatic-medicinal and culinary plants
		D13D3	Industrial plants-others
		D14A	Fresh vegetables, mellons, strawberries-outdoor-openfield
	D14-D15:		
	FRESH VEGETABLES, MELLONS, STRAWBERRIES	D14B	Fresh vegetables, mellons, strawberries-outdoor- market gardening
		D15	Fresh vegetables, mellons, strawberries under glass
	D16-D17:	D16	Flowers and ornamental plants outdoor
	FLOWER AND ORNAMENTAL PLANTS	D17	Flowers and ornamental plants under glass
		D18A	Forage plants-temporary grass
	D18: FORAGE PLANTS	D18B	Forage-plants-other green fodder-total
		D18B1	Forage-plants-other green fodder-green maize
		D18B2	Forage-plants-other green fodder-leguminous plants
		D18B3	Forage-plants-other green fodder-others
	D19-D20:	D19	Seeds and seedlings
	OTHER ARABLE CROPS	D20	Other crops
	D21: FALLOW LAND	D21	Fallow land
E: KITCHEN GARDENS	E: KITCHEN GARDENS	E	Kitchen gardens
F:	F:	F01	Permanent grassland and meadow-pasture and meadow
PERMANENT PASTURES AND MEADOWS	PERMANENT PASTURES AND MEADOWS	F02	Permanent grassland and meadow-rough grazing
G:	G1: FRUIT AND BERRY PLANTATIONS	GO1A	Fruit and berry plantations-temperate climate
PERMANENT CROPS		GO1B	Fruit and berry plantations-subtropical climate
		GO1C	Fruit and berry plantations-nuts
	G2: CITRUS PLANTATIONS	G02	Citrus plantations
	G3: OLIVE PLANTATIONS	GO3A	Olive plantations-table olives
		GO3B	Olive plantations-oil production
	G4: VINEYARDS	GO4A	Vineyards-quality wine
		GO4B	Vineyards-other wines
		GO4C	Vineyards-table grapes
		GO4D	Vineyards-raisins
		G5: NURSERIES	G05
	G6: OTHER PERMANENT CROPS	G06	Other permanent crops
	G7: PERMANENT CROPS UNDER GLASS	G07	Permanent crops under glass

H: OTHER LAND	H0103: NON-UTILIZED AGRICULTURAL LAND	H01	Unutilized agricultural land which is no longer farmed, for economic, social or other reasons
		H03	Other land occupied by buildings, pleasure gardens, etc.
	H02: WOODED AREA	H02	Woodland
1: COMBINE D AND SUCCESSIVE SECONDARY CROPPING, MUSHROOMS,	101: SUCCESSIVE SECONDARY CROPS	101A	Successive secondary crops-non fodder cereals
		101B	Successive secondary crops-non fodder pulses
		101C	Successive secondary crops-non fodder oil-seed plants
		101D	Successive secondary crops-others total
	102: MUSHROOMS	102	Mushrooms
	103: IRRIGATED AREA	103A	Total irrigable area
		103B	Irrigated once a year-total
	104: AREA COVERED BY GREENHOUSES IN USE	104	Area covered by greenhouses in use
	105: COMBINED CROPS	105A	Combined crops-agricultural-forestry
		105B	Combined crops-permanent-annual
105C		Combined crops-permanent-permanent	
105D		Combined crops-others	

**Table 2.** Linkage between the 2000 FSS and the new CLC nomenclatures in Greece

New CLC		FSS	
LEVEL 1	LEVEL 2	LEVEL 1	LEVEL 2
1. Artificial surfaces (Man-made areas)	1.1 Urban fabric (Build-up areas, urban agglomerations)		-
	1.2 Industrial and commercial units (Industrial or commercial zones)		-
	1.3 Transport units		-
	1.4 Mine, dump and construction sites (Mines, waste disposal sites)		-
	1.5 Artificial, non-agricultural vegetated areas sport and cultural activity sites (Artificial or non-agricultural green areas)		-
2. Agricultural areas	2.1 Arable land (Areas under arable crops)	Utilized agricultural areas D+G+E	D=D01+D02+D03+ D04+D05+ D06+D07+D08+D09+D10+ D11+D12+D13+D14+D15+ D16+D17+D18+D19+
	2.2 Permanent crops (Areas under permanent crops)		G=G01+G02+G03+G04+G05 +G06+G07
	2.3 Pastures (Areas under meadow or pasture)		F=F01+ F02
	2.4 Heterogeneous agricultural areas (Areas with mixed uses -mixed farmland)		105A+I05B
3. Forests and semi-natural areas	3.1 Forests (Forested areas)		H02: only the private forests
	3.2 Transitional woodland /shrub		H01: only the private uncultivated areas for economic, social or other reasons
	3.3 Shrub and/or herbaceous vegetation associations (Areas with mixed shrub/grassy vegetation)		
	3.4 Open spaces with little or no vegetation (Areas with little or no vegetation)		
4. Surfaces under water	4.1 Inland water		
	4.2 Inland wetlands		
	4.3 Coastal wetlands		

**Table 3.** Results showing the differences (%) in arable areas, areas under permanent crops and cultivated areas (D+E) as they recorded by the 2000 FSS and the new CLC nomenclatures.

Regions (NUTS II)	Districts (NUTS III)	Arable Areas (% difference)	Areas under Permanent Crops (% difference)	Cultivated Areas (% difference)
ANATOLIKI MAKEDONIA & THRAM	DRAMA	45	-93	42
	KAVALA	64	-45	31
	EVROS	24	44	25
	XANTHI	33	-67	32
	RODOPI	31	89	32
<b>TOTAL</b>		33	-27	30
KENTRIKI	IMATHIA	42	-91	-12
MAKEDONIA	SALONIKI	4	-49	3
	KILKIS	-7	-39	-7
	PELLA	-31	-77	-47
	PIERIA	-7	-79	-14
	SERRES	42	-81	37
	CHALKIDIKI	54	-9	34
<b>TOTAL</b>		15	-61	4
DYTIKI MAKEDONIA	GREVENA	20	-68	18
	KASTORIA	-21	-35	-22
	KOZANI	4	27	5
	FLORINA	-3	-44	-4
<b>TOTAL</b>		3	-14	2
<b>TOTAL MAKEDONIA</b>		18	-52	12
KRITI	IRAKLIO	-71	4	-4
	LASITHI	54	47	48
	RETHIMNO	-91	-7	-24
	CHANIA	-72	4	-4
<b>TOTAL</b>		-66	6	-3

**Table 4.** Results showing the actual values and the corresponding differences in the class of pastures and meadows as they recorded by the 2000 FSS and the new CLC nomenclatures.

	Districts	Pastures and meadows (ha)			
	(NUTS II)	(NUTS III)	2000 FSS	new CLC	Difference
ANATOLIKI	DRAMA		1,294	31,380	30,086
MAKEDONIA	KAVALA		760	19,810	19,050
	EVROS		4,353	13,870	9,517
	XANTHI		81	11,910	11,829
	RODOPI		1,733	13,520	11,787
TOTAL			8,221	90,490	82,269
KENTRIKI	IMATHIA		860	9,840	8,980
MAKEDONIA	SALONIKI		473	25,020	24,547
	KILKIS		5,310	40,680	35,370
	PELLA		2,458	25,910	23,452
	PIERIA		3	6,570	6,567
	SERRES		6,246	28,520	22,274
	CHALKIDIKI		2,780	5,330	2,550
TOTAL			18,130	141,870	123,740
DYTIKI	GREBENA		315	25,890	25,575
MAKEDONIA	KASTORIA		822	29,840	29,018
	KOZANI		794	70,610	69,816
	FLORINA		5,477	27,200	21,723
TOTAL			7,408	153,540	146,132
TOTAL MAKEDONIA			33,759	385,900	352,141
KRITI	IRAKLIO		36,412	69,070	32,658
	LASITHI		16,817	61,631	44,814
	RETHIMNO		62,470	53,241	-9,229
	CHANIA		63,410	40,167	-23,243
TOTAL			179,109	224,109	45,000



**Table 5.** Results showing the actual values and the difference (%) in the class of agricultural areas as they recorded by the FSS and the new CLC nomenclatures, It also shows the average parcel area.

Regions (NUTS II)	Districts (NUTS III)	Agricultural Areas (ha)		Agricultural Areas (% difference)	Average parcel area (ha)
		2000 FSS	New CLC		
ANATOLIKI MAKEDONIA	DRAMA	47,193	104,720	122	0.78
	KAVALA	44,860	92,390	106	0.55
	EVROS	150,252	231,060	54	0.64
	XANTHI	37,214	69,940	88	0.69
	RODOPI	74,941	121,230	62	0.62
	<b>TOTAL</b>		<b>354,460</b>	<b>619,340</b>	<b>75</b>
KENTRIKI MAKEDONIA	IMATHIA	53,894	95,690	78	0.82
	THESSALONIKI	129,483	222,840	72	0.79
	KILKIS	106,027	172,420	63	0.90
	PELLA	77,660	151,640	95	0.61
	PIERIA	45,543	74,950	65	0.75
	SERRES	144,947	234,670	62	0.58
	CHALKIDIKI	77,274	147,270	91	0.77
<b>TOTAL</b>		<b>634,828</b>	<b>1,099,480</b>	<b>73</b>	<b>0.72</b>
DYTIKI MAKEDONIA	GREVENA	41,432	93,810	126	0.80
	KASTORIA	24,887	74,260	198	0.58
	KOZANI	88,170	166,260	89	0.58
	FLORINA	52,952	90,960	72	0.55
<b>TOTAL</b>		<b>207,441</b>	<b>425,290</b>	<b>105</b>	<b>0.60</b>
<b>TOTAL MAKEDONIA</b>		<b>1,196,729</b>	<b>2,144,110</b>	<b>79</b>	<b>0.67</b>
KRITI	IRAKLIO	139,733	221,982	59	0.40
	LASITHI	37,864	127,252	236	0.44
	RETHIMNO	101,182	115,842	14	0.87
	CHANIA	109,191	116,472	7	0.83
<b>TOTAL</b>		<b>387,970</b>	<b>581,548</b>	<b>50</b>	<b>0.57</b>