

MATRICULATION AND SECONDARY EDUCATION CERTIFICATE **EXAMINATIONS BOARD**

INTERMEDIATE MATRICULATION LEVEL **2019 FIRST SESSION**

SUBJECT: Chemistry 22nd May 2019 DATE:

TIME: 9:00 a.m. to 12:05 p.m.

Useful information

Molar das constant $R = 8.31 \text{ Imol}^{-1}\text{K}^{-1}$

Relative atomic masses: $H=1$, $C=12$, $O=16$, $K=39$, $Ca=40$, $Fe=56$, $I=127$ A Periodic Table is included.
SECTION A
Answer ALL questions in this section.
 (a) A hydrocarbon Y contains 85.7% carbon and 14.3% hydrogen by mass. (i) Calculate the empirical formula of the hydrocarbon Y.
(1)
(ii) The molar mass of the hydrocarbon Y is 56 g. Deduce the molecular formula of the hydrocarbon Y.
(1)
(b) The analyst who wanted to identify the hydrocarbon Y, was certain that the compound was a saturated hydrocarbon. Suggest a structure for this saturated hydrocarbon Y. Name the hydrocarbon Y.
Structure:
Name: (1) (Total: 3 marks)

2.	The standard heat of formation of water in the gaseous state is -242 kJmol ⁻¹ . (a) Write an equation which represents the standard heat of formation of water in the gaseous state.
	(1)
	(b) Circle the correct value of the standard heat of combustion of gaseous hydrogen.
	-121 kJmol^{-1} -80.6 kJmol^{-1} -242 kJmol^{-1} -484 kJmol^{-1} (1)
	(c) Why is hydrogen considered to be a clean fuel?
	(1)
	(Total: 3 marks)
3.	There are four isomeric alkenes of molecular formula C_4H_8 . In the space below, draw the structure of each isomer. Write the name of each isomer under each structure.
4.	(Total: 4 marks) Carbon-14 in the atmosphere undergoes beta-decay and produces 60 atoms of nitrogen-14 every hour for each gramme of carbon. The disintegration rate is 60 counts hour ⁻¹ g ⁻¹ . (a) Write a balanced nuclear equation which shows the decay reaction of carbon-14.
	(b) A small piece of sea shell which was embedded in a rock was found to have a count of 3.75 counts hour -1g -1. The half-life of carbon-14 is 5730 years. Estimate the age of the seashell.
	(2) (Total: 4 marks)

5.	(a) Draw dot-cross diagrams for the following molecules:				
	Aluminium chloride (AICI	Ammonia (NH ₃)			
	Ammonium ion (NH ₄ ⁺)	Hydronium ion (H₃O⁺)			
	, ,	, , ,			
			(2)		
	(b) In the table below, w	rite the shape of each of the species.			
	Species	Shape			
ΑI	uminium chloride (AlCl ₃)				
	Ammonia (NH₃)				
	Ammonium ion (NH ₄ +)				
	Hydronium ion (H₃O ⁺)		(2)		
		(Total: 4 marks)		

Please turn the page.

6.	Polyvinyl chloride (PVC) is an addition polymer made from the monomer monochloroethene (a) What is an addition polymer?		
_			
	(1½)		
	(b) Draw the structural formula of monochloroethene.		
	(1)		
	(c) Draw the repeating unit of the polymer PVC.		
	(1½) (Total: 4 marks)		
7.	The reaction mechanism of the addition of hydrogen bromide to ethene is described as a		
	multistep reaction mechanism. (a) Draw a diagram to show the mechanism and the steps of this reaction. (In your diagram, use arrows to show the movement of electrons.)		
	(b) Explain the term rate-determining step.		
	(1)		
	(c) In your diagram in part 7(a), label clearly the rate determining step. (1) (Total: 4 marks)		

- 8. The hydrogenation of ethene is an exothermic reaction. This reaction is catalysed by Raney Nickel.
 - (a) Explain in terms of activation energy, what Raney Nickel does as a catalyst in this reaction.

____(1)

(b) Draw an energy profile diagram for the catalysed and the non-catalysed hydrogenation reaction. In your diagram label the activation energy of the catalysed reaction ($E_{a/cat}$), the activation energy of the non-catalysed reaction ($E_{a/non-cat}$) and the heat of the reaction (ΔH).

(Total: 4 marks)

Please turn the page.

SECTION B

Answer ALL questions in this section.

9.	(a)) The following two equations represent addition reactions in organic chemistry. They show the enthalpies of reaction of the hydrogenation of cyclohexene and the hydrogenation of benzene. Both compounds, when hydrogenated, give the same product.			
		Reaction 1			
		Reaction 2			
		(i) Describe the bonding between the atoms in the benzene molecule.			
			(1)		
		(ii) Explain why the enthalpy of hydrogenation in Reaction 2 is less exothermic three times the enthalpy of hydrogenation in Reaction 1.	c than		
			(1)		
	(b)	(i) Benzene undergoes substitution reactions rather than addition reactions. E this statement.	_ , ,		
			(1)		
		(ii) Give an equation for the mononitration of benzene.	_ , ,		
			(1)		
		(iii) Name the reagents which are used in the above reaction.	(1)		
		(iv) State the conditions which are used in this reaction.	(-)		
		(Total: 6 m	(1) narks)		

A mixture of gases composed of 44 g of propane and 224 g of oxygen is ignited in a vessel of volume 10 dm ³ . The temperature rises from 25 °C to 327 °C. (i) Find the number of moles of propane and the number of moles of oxygen present in the mixture.
 (2)
(ii) Write an equation for the combustion of propane.
 (½)
(iii) Find the total number of moles of gas left immediately after the reaction at 600 K.
 (1)
(iv) Calculate the pressure inside the vessel immediately after the reaction.
(1)
 The carbon dioxide which is produced in the combustion reaction is bubbled through lime water. The following equation represents this reaction. $ \text{Ca}(\text{OH})_2 \text{ (aq)} + \text{CO}_2 \text{ (g)} \longrightarrow \text{CaCO}_3 \text{ (s)} + \text{H}_2\text{O (I)} $ Calculate the mass of calcium carbonate which is precipitated.
 ($1lau_2$) (Total: 6 marks)
Sodium, aluminium and sulfur are elements in Period 3. Describe the reactivity (if any) of the oxides of these elements with water. (Include chemical equations where relevant.)
(3)

This question continues on the next page.

dioxide.

(b) The following table gives the melting point of three Period 3 oxides.

Oxide	Melting point (^o C)	
Na ₂ O	1132	
SiO ₂	1700	
SO ₂	- 72	

Explain the differences in the melting points of sodium oxide, silicon dioxide and sulfur

(3)
(Total: 6 marks)
. (a) Potassium iodate reacts with potassium iodide in acid solution to release iodine. The

12. (a) Potassium iodate reacts with potassium iodide in acid solution to release iodine. The redox ionic equation for this reaction is:

$$IO_3^-(aq) + 5I^-(aq) + 6H^+(aq) \longrightarrow 3I_2(aq) + 3H_2O(I)$$

(i) Give the oxidation number of iodine in:

Γ: ______ (½)

IO₃⁻: ______

(ii) Write a balanced ionic equation to show the oxidation half-reaction in the above reaction.

_____(1)

(iii) Write a balanced ionic equation to show the reduction half reaction in the above reaction.

_____(1)

(b) In an experiment, 0.4 g of potassium iodate (KIO_3) is dissolved in water and made up to 250 cm³ in a volumetric flask. Three 25 cm³ samples of the solution were separately added to an excess of potassium iodide in acid solution in conical flasks. The iodine which formed in each flask reacted with sodium thiosulfate from a burette in a titration.

$$I_2 + 2S_2O_3^{2-} \longrightarrow S_4O_6^{2-} + 2I^{-}$$

The following table gives the three titre values of the titrations.

Burette reading	Titration 1	Titration 2	Titration 3
Final (cm ³)	22.45	32.20	34.30
Initial (cm ³)	0.00	10.00	12.00

(i) Calculate the number of moles of potassium iodate in the 0.4 g sample.
(1½)
(ii) Calculate the number of moles of potassium iodate in the 25 cm ³ sample placed in each conical flask.
(½)
(iii) Calculate the number of moles of iodine which were formed in each conical flask.
(1½)
(iv) Calculate the number of moles of thiosulfate which reacted with iodine in each conical flask.
(½)
(v) What is the concentration of the sodium thiosulfate solution?
(1) (Total: 6 marks)
13. Ethene is used as a starting material in the industrial production of ethanol. The ethene is reacted with steam at around $6 \times 10^6 \text{ Nm}^{-2}$ and at a temperature of 300 °C. The catalyst in this reaction is phosphoric acid. Under these conditions, 5% of the reactants are converted into ethanol. $CH_2 = CH_2 (g) + H_2O (g) \qquad CH_3CH_2OH (g) \qquad \Delta H^e = -46 \text{ kJmol}^{-1}$
(a) (i) Explain, in terms of Le Chatelier's principle, why a high pressure favours the production of ethanol.
(2)
(ii) Give ONE disadvantage of using a high pressure in an industrial process.
(1½)

This question continues on the next page.

	(i)	Explain, in terms of Le Chatelier's principle, why a low temperature favours the formation of ethanol.
		(2)
	(ii)	Give ONE disadvantage of using a low temperature in an industrial process.
		(1/2)
(c)	(i)	How does phosphoric acid affect the position of equilibrium in the production of ethanol?
		(1/2)
	(ii)	What happens if phosphoric acid is not used in the reaction?
SECTIO	ON C	($^{(1/2)}$
Answe of this		TWO questions from this section. Write your answers on the lined pages let.
of this	there (i) (ii) (iii) (iv) (v) (vi) (vii)	e are acids, bases and alkalis. Give the Arrhenius definition of the terms acid and base, i.e. in terms of H ⁺ and OH ⁻ .(2) Distinguish between a base and an alkali. There are strong and weak acids and bases. Distinguish between the terms strong and weak in this context. Give an example of each of the following: a strong acid, a weak acid, a strong base, a weak base. Explain the term amphoteric. Water is an amphoteric compound. Give TWO reactions of water to show this behaviour. Give the Lowry-Bronsted definition of the terms acid and base. Nitric acid reacts with methanol as follows:
of this	there (i) (ii) (iii) (iv) (v) (vi) (vii)	e are acids, bases and alkalis. Give the Arrhenius definition of the terms acid and base, i.e. in terms of H ⁺ and OH ⁻ .(2) Distinguish between a base and an alkali. There are strong and weak acids and bases. Distinguish between the terms strong and weak in this context. Give an example of each of the following: a strong acid, a weak acid, a strong base, a weak base. Explain the term amphoteric. Water is an amphoteric compound. Give TWO reactions of water to show this behaviour. Give the Lowry-Bronsted definition of the terms acid and base.

- 15. (a) Potassium manganate(VII), or potassium permanganate, KMnO₄, in acidified solution, is titrated against a solution containing iron(II) ions, and turns to manganese(II) ions.
 - (i) Write the half equation that shows the change from acidified manganate(VII) to manganese(II) ions. (2)
 - (ii) Write the half equation that shows the change that iron(II) undergoes when it is titrated with acidified manganate(VII). (1)
 - (iii) Join the **TWO** half equations in parts (i) and (ii), and write the full redox reaction between acidified manganate(VII) and iron(II). (1)
 - (iv) Why is there no need for an indicator in the titration between acidified manganate(VII) and iron(II)? (1)
 - (b) The following procedure was carried out:
 - An iron tablet was weighed and transferred to a beaker; 100 cm³ of deionised water were added, and heated to dissolve the tablet.
 - The solution was allowed to cool and transferred to a 250 cm³ volumetric flask, and made up to the mark with deionised water.
 - 1000 cm³ of 0.01 moldm⁻³ potassium manganate(VII) solution were prepared; it was acidified by 2 moldm⁻³ sulphuric acid.
 - 10 cm³ of the acidified 0.01 moldm⁻³ potassium manganate(VII) solution were pipetted into a conical flask.
 - The unknown iron(II) solution was run into the flask from a burette, recording the end point.
 - The titration was repeated until concordant results are obtained. The results were:

Titre	Initial (cm ³)	Final (cm ³)	Titre value (cm ³)
1	0.00	22.08	
2	0.05	27.95	
3	0.00	27.90	

- (i) Why was the solution allowed to cool before it was transferred to the 250 cm³ volumetric flask? (1)
- (ii) What is meant by concordant results? Indicate the concordant titre values. (1)
- (iii) Find the mass of iron per tablet in mg.
- (c) A dry sample of a mixture of $CaCO_3$ and $CaCl_2$ has a mass of 1.45 g. It was dissolved in 25.00 cm³ of 0.98 moldm⁻³ HCl solution. A volume of 21.48 cm³ of 0.09 moldm⁻³ NaOH solution was used to titrate the excess HCl.

What was the percentage of CaCl₂ in the original mixture?

(Total: 20 marks)

- 16. (a) Chlorine (and bromine) react with and oxidise thiosulfate. Iodine oxidises thiosulfate too, but it does so differently.
 - (i) Give the reaction of chlorine with thiosulfate.

(2) (2)

(6)

- (ii) Give the reaction of iodine with thiosulfate.
- (iii) Considering the reaction of chlorine with thiosulfate, by assigning oxidation numbers to every element, indicate what is oxidised and what is reduced. (5)
- (iv) Considering the reaction of iodine with thiosulfate, by assigning oxidation numbers to every element, indicate the oxidising agent and the reducing agent. (5)
- (b) We speak of the relative oxidising power of the halogens. Explain, by considering the **THREE** halogens chlorine, bromine and iodine. Include chemical equations wherever necessary. (6)

(Total: 20 marks)

Please turn the page.

17.	Explain each of the following statements, illustrating your answer with suitable chemical reactions wherever possible.	al
	(a) Esters are produced through one of the typical reactions of alcohols, in the presence of	of 4)
		4)
	(c) Primary, secondary and tertiary alcohols can be distinguished through their oxidation	
		5)
		5)
	(Total: 20 marks	
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_
		_

DO NOT WRITE ABOVE THIS LINE

DO NOT WRITE ABOVE THIS LINE

PERIODIC TABLE

1											_			_	_		_			
Name	VIII	4 He	0,0	2 2	10	40	Ar	18	84	Ϋ́	36	131	Xe	54	222	&	98			
Name	VII		10	J D	4 6	35.5	Ü	17	80	Br	35	127	I	53	210	At	85			
Name	VI		16	2 0	> ∞	32	S	16	19	Se	34	128	Le	52	209	Po	84			
Scalative A Atomic X Atomic Ato	>		7	± 7	ζ Γ	31	4	15	75	As	33	122	Sb	51	209	Bi	83			
$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	IV		5	1 (ہ ر	28	Si	14	73	g	32	119	Sn	50	207	Pb	82			
Sc Ti V Cr Min Fe Co Ni Cu			=	= p	a 50	27	A	13	70	Ga	31	115	П	46	204	I	81			
Sc Ti V Cr Min Fe Co Ni			_						65	Zn	30	112	Cq	48	201	Hg	80			
Sc Ti V Cr Min Fe Co									63.5	Cn	29	108	Ag	47	197	Au	19			
Key 45 48 51 52 55 56 Sc Ti V Cr Mn Fe 21 22 23 24 25 56 89 91 93 96 99 101 Y Zr Nb Mo Tc Ru 39 40 41 42 43 44 139 178.5 181 184 186 190 La Hf Ta W Re Os 57 72 73 74 75 76 89 89									59	Z	28	106	Pd	46	195	F	78			
Ac		Atomic	-Number						59	ပိ	27	103	Rh	45	192	H	77			
89 91 93 96 W	Key	< ⋈							99	Fe	56	101	Ru	44	190	SO	9/			
45 48 51 Sc Ti V 21 22 23 89 91 93 Y Zr Nb 39 40 41 139 178.5 181 La Hf Ta 57 72 73 Ac		Relative atomic							55	Mn	25	66	Te	43	186	Re	75			
45 48 48 Sc Ti 22 22 22 39 91 40 139 178.5 La Hf 57 72 Ac 89									52	ڻ	24	96	Mo	42	184	A	74			
Sc 227									51	>	23	93	Z	41	181	La	73			
									48	Ξ	22	91	Zr	40	178.5	Hf	72			
11 Be 9 9 9 9 12 24 4 4 6 6 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8									45	Sc	21	68	×	39	139	La	57	227	Ac	68
	П			۷ ۾	Pe 4	24	Mg	12,	40	Ca	70	88	Sr	38	137	Ba	99	226	Ra	88
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	I	- н -	, ,	\ ;	n ا	23	Na	. 11	39	K	19	85	Rb	37	133	C	55	223	Fr	87

175	Lu	71	. 260	Lr	103
173	ΛP	70	259	No No	102
169	Tm	69	258	Md	101
191	Er	89	257	Fm	100
165	Ho	29	252	Es	66
162	Dy	99	251	Ct	86
159	·Tb	9	247	Bk	97
157	Вd	64	247	Cm	96
152	Eu	63	243	Am	95
150	Sm	62	244	Pu	94
147	Pm	19	237	ď	93
144	PN	09	238	n	92
141	Pr	59	231	Pa	91
140	ů	28	232	Th	90